PUPT–1324
hepth@xxx/9205100

arXiv:hep-th/9205100v2 25 Aug 1992

A note on the 3D Ising model as a string theory*

Jacques Distler
Joseph Henry Laboratories
Princeton University
Princeton, NJ 08544 USA

It has long been argued that the continuum limit of the 3D Ising model is equivalent to a string theory. Unfortunately, in the usual starting point for this equivalence – a certain lattice theory of surfaces – it is not at all obvious how to take the continuum limit. In this note, I reformulate the lattice theory of surfaces in a fashion such that the continuum limit is straightforward. I go on to discuss how this new formulation may overcome some fundamental objections to the notion that the Ising model is equivalent to a string theory. In an appendix, I also discuss some aspects of fermion doubling, and the lattice fermion formulation of the 2D Ising model.

5/92

* Email: distler@puhep1.princeton.edu .
Research supported by NSF grant PHY90-21984.

1. Introduction

The 2D Ising model was first solved by Onsager [1] in the 1940's, and has been reformulated and re-solved by many others subsequently (some references relevant to my discussion are [2,3,4,5,6,7]). It has been a rich and fruitful source of insights into 2D critical phenomena.

The 3D Ising model has, so far, resisted an exact solution. Though much is known about its critical behaviour, both analytically and numerically, many have hoped that it might yet yield, if not to an exact solution, then at least to a closed-form calculation of its critical exponents. One of the most elegant suggestions for reformulating the 3D Ising model has been to try to recast it as a string theory, that is, as a theory of surfaces immersed in 3 dimensions.

Despite much work on this subject [8,9,10,12,13], not much progress has been made. In this paper, I will reconsider the traditional approach to recasting the 3D Ising model as a string theory and try to suggest where it runs into difficulties. I will then suggest an alternate approach which has the promise of overcoming these difficulties.

Anyone who tries to to convince you that the continuum limit of the 3D Ising model is equivalent to some string theory had better be prepared to address at least two key objections:

- String theory would seem to have far too many degrees of freedom to be equivalent to what is, after all, a *field theory* in 3 dimensions.
- Intimately related [14] to this large density of states is the fact that string theory generically has a tachyon in its spectrum, which is surely *not* a property of the 3D Ising model. In practice, the tachyon leads to a divergence of the string integrand near surfaces with long thin tubes. In other words, these configurations tend to dominate the functional integral, whereas we expect that near the critical point of the Ising model, domains of all sizes become important.

These problems are *very generic* and are more or less independent of the world-sheet theory (so long as it has a large enough number of degrees of freedom). Since precisely the same problems are currently faced by noncritical string theory (for $D > 1$), it is very interesting to see how the 3D Ising model manages to solve them. This is the main motivation behind the present work.

The plan of this paper is as follows. In section 2, I review the 2D Ising model from a point of view which will generalize to 3 dimensions. In section 3, I discuss the 3D Ising

model. I discuss the approach adopted in previous attempts to recast it as a string theory and suggest an alternate approach, where taking the continuum limit is more straightforward. In section 4, I return to the problems just mentioned, and discuss how, in this new approach to the 3D Ising model, they might be solved.

There are three appendices. Appendix A is devoted to the lattice fermion representation of the 2D Ising model, and how it avoids fermion doubling. Appendix B contains a brief speculation on the subject of the equivalence between the Nambu-Goto and Polyakov strings in 3 dimensions. Appendix C is a review of the definition of fermions on nonorientable manifolds.

2. Review of the 2-D Ising model

Let us start our discussion by recalling the situation in the 2-D Ising model. The partition function is obtained by summing over all configurations of spins $\sigma_i = \pm 1$ on the sites of the lattice

$$\mathcal{Z}[\beta] = \sum_{\{\sigma\}} \mathrm{e}^{-\beta \sum_{<ij>} (1-\sigma_i \cdot \sigma_j)} \quad . \tag{2.1}$$

The action $S = \sum_{<ij>}(1 - \sigma_i \cdot \sigma_j)$ only receives contributions from links across which the neighbouring spins are anti-aligned. If we have a domain of up spins ($\sigma = +1$) next to a domain of down spins ($\sigma = -1$), the contribution to the action is proportional to the length of the boundary separating the two domains. Thus we can contemplate replacing (2.1) by a sum over *curves* (on the dual lattice) weighted by their lengths [3]

$$\mathcal{Z}[\beta] = 2Z[\beta] \stackrel{?}{=} 2 \sum_{\{\gamma\}} \mathrm{e}^{-2\beta L[\gamma]} \tag{2.2}$$

The factor of 2 in this formula simply accounts for the $\mathbb{Z}_2$ symmetry of the Ising model, in which flipping all of the spins $\sigma_i \to -\sigma_i$ produces the same configuration of curves. To make this formula correct, we should first of all restrict ourselves to closed (but not necessarily connected) curves γ which traverse each link of the (dual) lattice only *once* (or zero times). This restriction to so-called *unicursal* curves will be relaxed later, but for the moment it is still not enough to make the equality (2.2) correct.

The problem is that the curve γ can self-intersect. If we consider the configuration of 4 spins shown in Fig. 1a, we see that the corresponding curve γ has four links which apparently meet at a point. There are three ways to resolve this singular point (Fig. 1b,c,d).

That is to say, there are three different immersed curves whose images involve the same four links. Counting immersed curves as in (2.2), we would count each of these configurations separately. This is *overcounting* since they all correspond to the same configuration of spins. We can fix this by introducing the intersection number $n(\gamma)$ which counts the number of intersections of the immersed curve γ. Fig. 1d contributes 1 to $n(\gamma)$, while Fig. 1b,c do not intersect, and so contribute 0 to $n(\gamma)$. Instead of (2.2), consider

$$\mathcal{Z}[\beta] = 2Z[\beta] = 2\sum_{\{\gamma\}}(-1)^{n(\gamma)}\mathrm{e}^{-2\beta L[\gamma]} \quad . \tag{2.3}$$

This counts Fig. 1d with a minus sign relative to Fig. 1b,c. Two of the three configurations cancel leaving a *net* of one configuration, which agrees with the sum over spins.



Fig. 1: A configuration of spins, and three different immersed curves corresponding to it

Having introduced the factor of $(-1)^{n(\gamma)}$, we can relax the requirement that the curve γ be unicursal. This is good because the unicursal restriction is an extremely nonlocal one, and would like to replace it with something local. Let us define an *admissible* curve to be a closed (not necessarily connected) curve on the lattice which does not retrace itself. That is, no link is traversed twice in *successive steps*. The three configurations Fig. 2a,b,c are allowed; the configuration Fig. 2d is not allowed.



Fig. 2: Three admissible configurations and one inadmissible one



Fig. 3: An admissible configuration and its resolutions.



Fig. 4: The admissible curve γ which contributes to the partition function.

Consider the admissible configuration in Fig. 3a. The central link is occupied twice. This configuration corresponds to four different admissible curves (Fig. 3b,c,d,e). But with the sign factor in (2.3), two of these configurations get counted with a plus sign and two with a minus. Thus they cancel, leaving only (Fig. 4) as the curve γ corresponding to this configuration of spins.

In fact, we claim

$$\sum_{\{\substack{unicursal \\ \gamma}\}} (-1)^{n(\gamma)} \mathrm{e}^{-2\beta L[\gamma]} = \sum_{\{\substack{admissible \\ \gamma}\}} (-1)^{n(\gamma)} \mathrm{e}^{-2\beta L[\gamma]} \tag{2.4}$$

To prove this, we need to show that the cancellation seen in the above example is a general phenomenon whenever any link is multiply-occupied. Consider a vertex at which four links, occupied k_1 through k_4 times respectively, come together (Fig. 5a). Let k_1 be the largest of the k_i.

The easy case to handle is when $k_1 = k_2 + k_3 + k_4$. In this case, each line from links $2, 3, 4$ *must* be connected to a line from link 1. The number of different ways of doing this is given by the number of permutations of the k_1 lines on link 1. The sign factor associated to a particular way of connecting the lines is given by the *signature* of the corresponding permutation,

$$(-1)^n = \sigma(g), \qquad g \in S_{k_1} \quad .$$



Fig. 5: A vertex at which four multiply-occupied links meet

Summing over all possible ways of connecting the lines amounts to summing over S_{k_1} the group of permutations of k_1 objects. But it is elementary that

$$\sum_{g \in S_{k_1}} \sigma(g) = \begin{cases} 0 & k_1 > 1 \\ 1 & k_1 = 1 \ . \end{cases}$$

So we get zero unless $k_1 = 1$, in which case, one of $k_{2,3,4} = 1$, and the other two are zero.

The harder case is when $k_1 + 2p = k_2 + k_3 + k_4$. In this case, we connect up $2p$ lines among $k_{2,3,4}$, leaving $k_2 + k_3 + k_4 - 2p$ lines to connect to k_1. (Later, we will sum over the different ways of making the first p connections.) But this reduces the computation of the sign factor to an overall sign (determined by our choice of the first p connections) multiplied by the same $\sigma(g)$ (see Fig. 5b). Again, summing over all $g \in S_{k_1}$ gives zero unless $k_1 = 1$.

This argument, a much-simplified version of the one appearing in [4], allows us to replace unicursal curves γ by admissible curves in (2.3),

$$Z[\beta] = \sum_{\left\{ \substack{admissible \\ \gamma} \right\}} (-1)^{n(\gamma)} \mathrm{e}^{-2\beta L[\gamma]} = \exp\left[\sum_{\left\{ \substack{connected \\ admissible \\ \gamma} \right\}} (-1)^{n(\gamma)} \mathrm{e}^{-2\beta L[\gamma]} \right] \quad . \qquad (2.5)$$

Note that the sum over admissible curves has another advantage – it exponentiates – as a sum over unicursal curves would not.

From here it is relatively easy to see that the continuum limit of this theory is a theory of free majorana fermions. The most direct way is to define a theory of lattice fermions [6,7] whose partition function is reproduced by the same diagrammatic expansion as (2.3). Some issues related to this approach are discussed in Appendix A. More intuitively, we recognize (2.5) as the (Euclidean) partition function for a 1+1 dimensional quantum field theory, in a first-quantized formalism (familiar to any string theorist). The particles do not

interact, but they do have statistics. Every time two particle trajectories cross (the only way particles can be exchanged in 1+1 dimensions) we get a minus sign – so the particles are fermions.

The connection with free fermions becomes even more clear when we consider the theory on a lattice with periodic boundary conditions [5]. On a finite lattice with periodic boundary conditions, the sum over curves still overcounts configurations. Consider a loop which winds around one of the homology cycles of the torus. This doesn't correspond to any configuration of spins. More generally, γ can have total winding number p_t in the horizontal direction and q_t in the vertical direction. Only (p_t, q_t) *even* correspond to physically realizable spin configurations.

The solution is simple. First introduce "independent" horizontal and vertical couplings β_h, β_v. Define

$$Z(\beta_h, \beta_v) = \sum_{\{\gamma\}} (-1)^{n(\gamma)} \mathrm{e}^{-2(\beta_h L_h[\gamma] + \beta_v L_v[\gamma])} \tag{2.6}$$

where $L_{h,v}[\gamma]$ is the number of horizontal (vertical) links in γ ($L[\gamma] = L_h + L_v$). Our old partition function is just $Z[\beta] = Z(\beta, \beta)$. But now consider

$$Z(\beta + i\pi/2, \beta) = \sum_{\{\gamma\}} (-1)^{n(\gamma)} \mathrm{e}^{-2\beta L[\gamma]} (-1)^{p_t(\gamma) N_h}$$

where $N_{h,v}$ = horizontal (vertical) length of the lattice ($N = N_h N_v$). Similarly,

$$Z(\beta, \beta + i\pi/2) = \sum_{\{\gamma\}} (-1)^{n(\gamma)} \mathrm{e}^{-2\beta L[\gamma]} (-1)^{q_t(\gamma) N_v} \quad .$$

To reproduce the Ising partition function on a periodic lattice, we simply need to sum

$$\begin{aligned} \mathcal{Z}[\beta] &= \tfrac{1}{2}[Z(\beta, \beta) + Z(\beta + i\pi/2, \beta) \\ &\qquad + Z(\beta, \beta + i\pi/2) + Z(\beta + i\pi/2, \beta + i\pi/2)] \qquad (2.7a) \\ &= 2 \sum_{\{\gamma\}} \tfrac{1}{2}\Big(1 + (-1)^{p_t(\gamma) N_h}\Big) \tfrac{1}{2}\Big(1 + (-1)^{p_t(\gamma) N_h}\Big) (-1)^{n(\gamma)} \mathrm{e}^{-2\beta L[\gamma]} \qquad (2.7b) \end{aligned}$$

Following [5], we assume the $N_{h,v}$ are both *odd*, and we see that (2.7*b*) projects onto q_t, p_t both even. But the terms in the sum (2.7*a*) are clearly recognizable* as the sum over spin structures of the fermion. (This connection is made explicit in Appendix A). At the critical point, the contribution of the odd spin structure, $Z(\beta, \beta)$, vanishes.

* For a dirac fermion, one can follow [15] and represent the different spin structures by coupling the fermion to a flat $U(1)$ bundle. For a majorana fermion on the lattice, the same trick is available, provided we take the holonomy across each link of the lattice to be ± 1.

3. On to 3-D ...

The lesson that can be abstracted from the previous section can be summarized as follows. We wanted to recast the 2-D Ising model as a theory of immersed curves. However, if we are not careful, the sum over immersed curves overcounts configurations. The point is that there are two natural topologies that one can associate with a curve immersed in $\mathbb{R}^2$. One is the intrinsic topology of the curve. The other is the topology inherited from $\mathbb{R}^2$, the *extrinsic* topology. These are in general different. Points which are far apart in the intrinsic topology may be close together in the extrinsic topology. What is more, there are many possible intrinsic topologies which give rise to the same extrinsic topology. But the Ising model action depends only on the the extrinsic geometry of the immersed curve, and is insensitive to the intrinsic topology. Thus we needed to introduce a topological term $(-1)^{n(\gamma)}$ which distinguished between the different intrinsic topologies corresponding to a given extrinsic geometry and introduced cancellations between them.

In three dimensions, the boundaries between domains are closed surfaces, not curves. Again, we restrict ourselves at first to unicursal surfaces – those that occupy each plaquette (of the dual lattice) at most once. We wish to rewrite the Ising partition function as a sum over surfaces weighted by their area

$$\mathcal{Z}[\beta] = 2Z[\beta] \stackrel{?}{=} 2\sum_{\{\Sigma\}} \mathrm{e}^{-2\beta A[\Sigma]} \tag{3.1}$$

but, again, this overcounts configurations because the surface may have many intrinsic topologies corresponding to a given extrinsic topology.

One possible solution, suggested by Fradkin, Srednicki and Susskind [8] and elaborated by Polyakov and Dotsenko [11] and others [13], is a simple generalization of (2.3) to 3 dimensions,

$$\mathcal{Z}[\beta] = 2Z[\beta] = 2\sum_{\{\Sigma\}} (-1)^{L_{int}[\Sigma]} \mathrm{e}^{-2\beta A[\Sigma]} \tag{3.2}$$

where L_{int} is the length of the line of self-intersection of the surface Σ in lattice units. The basic idea can be seen in Fig. 6. Here we have three configurations (drawn as continuous surfaces), in which four sheets of the surface come together along a line. One possible intrinsic topology corresponding to this extrinsic topology is depicted in Fig. 6a. The surface is taken to self-intersect along the whole line. The top sheet on one side is connected to the bottom sheet on the other side, and vice versa. The other two surface differ from the first only in how they are connected across one (the left-most) link. In Fig. 6b, the top

sheet on one side is connected to the top sheet on the other. In Fig. 6c, it is connected to the bottom sheet on the same side. In each case, I've drawn a dashed curve to indicate the path taken by a "bug" which starts out on the top sheet above. Note that from the point of view of the bug (*i.e.* in the intrinsic topology) nothing special happens as it crosses the line of self-intersection.



Fig. 6: Three configurations, two of which must cancel in the sum over surfaces

Since the surface Fig. 6a self-intersects along one more link of the lattice than Fig. 6b,c, by the above rule it gets counted with a relative minus sign in the sum over surfaces (3.2).

It was originally believed that the continuum limit of (3.2) was the NSR string in 3 dimensions [9], and although Polyakov has subsequently suggested other possibilities for what the continuum theory might be, nothing is conclusively proven. Part of the difficulty is that sign factor $(-1)^{L_{int}[\Sigma]}$ oscillates very rapidly on the length scale of the lattice spacing. It is therefore not obvious how one is supposed to take the continuum limit. Also, the generalization from 2 to 3 dimensions of the argument that leads to fermions is not as straightforward as one might think[16] and it is not at all clear that (3.2) is equivalent to the NSR string.

One of the features that makes theories of surfaces in 3 dimensions so much richer than that of curves in 2 dimensions is that the possibilities for the intrinsic topologies are much more varied. In two dimensions, a curve γ was, in terms of its intrinsic topology, simply a disjoint union of circles. The only feature which distinguished different intrinsic topologies was the number of connected components. Surfaces have a much richer topological classification and we will make use of this fact to replace the sign factor $(-1)^{L_{int}[\Sigma]}$ with one that has a more obvious continuum limit.

Look again at Fig. 6. One thing is immediately evident is that one of the surfaces that we have drawn (Fig. 6b) differs in its intrinsic topology from the other two. Indeed, Fig. 6a,c are homotopic (they can be deformed continuously into one another). To see exactly how Fig. 6b is different, we need to make a short digression on the topology of 2-manifolds.

The topological classification of compact, connected 2-manifolds [17] is completely given by the cohomology of the manifold with $\mathbb{Z}_2$ coefficients. The basic theorem is that one can choose a basis for $\mathrm{H}^1(\Sigma, \mathbb{Z}_2)$ such that the generators have one of the following three intersection forms

$$\begin{array}{llll} \text{i)} & \alpha_i \cup \beta_j = \delta_{ij} & \text{ii)} \quad \alpha_i \cup \beta_j = \delta_{ij} & \text{iii)} \quad \alpha_i \cup \beta_j = \delta_{ij} \\ & & \gamma \cup \gamma = 1 & \gamma \cup \gamma = 1 \\ & & & \gamma \cup \tilde{\gamma} = 1 \end{array} \tag{3.3}$$

with all other cup products being zero. The indices i, j to run from 1 to g, where g is the number of "handles". The dimensions of $\mathrm{H}^1(\Sigma, \mathbb{Z}_2)$ in the three cases are, respectively, $2g$, $2g+1$, and $2g+2$. For $g = 0$, these correspond to the sphere, the projective plane, and the Klein bottle.

The Stiefel-Whitney classes of Σ are easily computed from Wu's formula [18], and the above cohomology ring:

$$\begin{array}{llll} \text{i)} & w_1 = 0 & \text{ii)} \quad w_1 = \gamma & \text{iii)} \quad w_1 = \tilde{\gamma} \\ & w_2 = 0 & w_2 = 1 & w_2 = 0 \end{array} \tag{3.4}$$

The first Stiefel-Whitney class, w_1 is the obstruction to orientability of Σ. The second Stiefel-Whitney class $w_2(\Sigma)$ is the obstruction [19] to defining fermions (or, more technically, to defining a *pin*-structure [20]) on Σ. For closed surfaces, it is the reduction modulo 2 of the Euler characteristic

$$w_2(\Sigma) = \chi(\Sigma) \quad \text{modulo } 2 \quad .$$

It is easy to see that the surface Fig. 6b has Euler characteristic (or equivalently w_2) which differs by 1 from that of Fig. 6a or c. In particular, Fig. 6b is non-orientable. To see this, consider the path of a "bug" in Fig. 7. As the bug traverses the dotted path, it



Fig. 7: A cycle on a non-orientable surface

comes back to itself with a reversal of orientation. The dotted path is essentially dual to the new $\mathbb{Z}_2$ cohomology cycle γ introduced above.

Let us try to make use of this difference in intrinsic topology to introduce the sign factors which we need. Instead of (3.2), let us take

$$\mathcal{Z}[\beta] = 2Z[\beta] = 2\sum_{\{\Sigma\}}(-1)^{w_2(\Sigma)}\mathrm{e}^{-2\beta A[\Sigma]} \tag{3.5}$$

Actually, when one wants to discuss disorder operators, *i.e.* when one wants to include surfaces Σ with boundaries, there are further sign factors which must be included:

$$Z[\beta, C = \partial\Sigma] = \sum_{\{\Sigma\}}(-1)^{w_1(C)+w_2(\Sigma)}\mathrm{e}^{-2\beta A[\Sigma]} \tag{3.6}$$

but we will, for the purposes of this paper, mostly restrict our attention to closed surfaces.

Note that non-orientable surfaces are *inevitable* if we want to play this game of cancelling surfaces with the same extrinsic geometry against each other. If we did not include Fig. 6b (which happens to be non-orientable), we would have had no way of cancelling Fig. 6a,c to leave a net of one configuration. This was just as true of the old weighting scheme (3.2) as for the new one (3.5). The only difference is that now we are trying to make direct use of the fact.

As an aside, let me point out an apparent paradox. As we have argued, (3.2) necessarily involves surfaces, like the projective plane, $\mathbb{R}\mathrm{P}^2$, which have nonvanishing w_2. This means that there is an obstruction to (the standard definition of) fermions on such surfaces. We can understand this obstruction physically in the simple case of $\mathbb{R}\mathrm{P}^2$. $\mathbb{R}\mathrm{P}^2$ can be built by gluing together a disk and a mœbius band along their common boundary (an

S^1). We can imagine doing the fermion path integral on $\mathbb{R}\mathrm{P}^2$ in stages: compute the path integral on the disk to obtain a state on the boundary S^1. Similarly, do the path integral on the mœbius band to obtain a state on its boundary. Finally, we obtain the path integral on $\mathbb{R}\mathrm{P}^2$ by taking the inner product of the state associated to the disk and the state associated to the mœbius band. It is easy to see that the state associated to the disk is in the NS sector (the nontrivial spin structure on S^1), whereas the state associated to the mœbius band is in the R sector (the trivial spin structure on S^1). Since these are orthogonal, the only consistent definition of the path integral is to take it to vanish identically. (If, however, we allow insertions, and consider amplitudes involving the insertion of an *odd* number of spin operators, we *can* obtain a nonzero result.) But this is not the behaviour one expects to find in the NSR string, where one expects $\mathbb{R}\mathrm{P}^2$ to have a dilaton tadpole [21], and it is certainly not the behaviour we want in the Ising model, where surfaces like $\mathbb{R}\mathrm{P}^2$ are supposed to cancel the contributions of other surfaces.

The resolution to this paradox is that on nonorientable manifolds, there are two *inequivalent* definitions of fermions. The "standard" definition indeed has as its obstruction w_2. The other definition, in turn, has as its obstruction, $w_2 + w_1 \cup w_1$. As can be seen from (3.3),(3.4), the latter obstruction vanishes identically on any 2-manifold. Clearly, the thing to do is to define the fermions in the nonorientable NSR string using the second definition. The existence of these two inequivalent definitions of fermions, though a "known" result [22], does not seem to be widely appreciated, even among open string theorists. I have therefore included a short summary in Appendix C. In any case, it is unlikely, even using the second definition of fermions, that (3.2) is related to the NSR string.

But let us return to our discussion of (3.5). Superficially, this seems to introduce cancellations of the desired sort, but does it work in detail? Does the lattice sum over surfaces precisely reproduce the Ising model sum over spins? In answering this question, we must first face up to the fact that we actually need to specify a *rule* for reconstructing intrinsic surfaces from a collection of plaquettes. This rule must have two crucial properties:

- It must be *local*. That is to say that the choice of how we connect together plaquettes at one link of the lattice cannot depend on distant information (say, on choices made at distant links.)
- It must be *intrinsic*. That is, if we have four sheets coming together at a link (Fig. 8), we can connect sheet 1 to sheet 2 and sheet 3 to 4, or we can connect 1 to 4 and 2 to 3, or we can connect 1 to 3 and 2 to 4. An intrinsic rule is one which is invariant under permutation of the labels of the sheets.



Fig. 8: Four sheets coming together on a line, and our notation for how to connect them

For later notational convenience, we will denote the three ways of connecting the sheets together by using a solid, dashed, or gray line to colour the link in question (Fig. 8).

In the 2D case, there was only one possible intrinsic, local rule for turning a collection of links into an immersed curve. We didn't even contemplate the existence of possible alternatives*. In the 3D case, we must face up to the existence of a choice.

From the continuum point of view, the lattice is providing a short-distance regulator for the theory. Different "rules" correspond to slightly different lattice regulators for the theory of surfaces. The notion of universality is simply that the details of the lattice regulator are irrelevant in the continuum limit. Because of the tachyon divergence, as we shall discuss in the next section, such a naive application of the principle of universality needs to be checked carefully.

Before I state the rule which we shall use, let me motivate it heuristically. The lattice acts as a regulator for the theory of surfaces basically by providing a minimal length for cycles on Σ. We want the cutoff to be *shorter* for cycles which carry the information about the orientability of Σ (*e.g.* the cycle γ in Fig. 7). As we have seen, these are cycles γ with

$$\gamma \cap \gamma = 1 \quad . \tag{3.7}$$

We will impose the condition that such cycles have length ≥ 1 in lattice units, whereas all other cycles must have length ≥ 2. Note that, because of the $\mathbb{Z}_2$ grading, it is consistent

* Note that if we were willing to forego the requirements of locality and intrinsicness, there would clearly exist rules which would eliminate the multiple counting – without the need to introduce the $(-1)^{n(\gamma)}$ factor.

to impose this shorter cutoff on cycles γ satisfying (3.7) (the sum of two such cycles has length ≥ 2, as it must). The choice of $1, 2$ lattice units is somewhat arbitrary, and I could double both of them, if you prefer, at the cost of somewhat complicating the statement of the rule to be given presently.

The rule for what you are allowed to do at one link where four sheets of Σ come together will depend only on how the sheets are connected immediately on either side of the link in question. That is to say, it will depend on what is happening at "nearest neighbour" links, but *not* on more distant information. It is therefore a *local* rule. The allowed configurations are depicted in Fig. 9. The circles represent how the sheets are connected on either side of the link in question (using the "colour" mnemonic introduced in Fig. 8). Notice that, in some of the allowed configurations, the way the sheets are connected changes at *half*-lattice spacings. As I said above, this can be avoided at the cost of somewhat complicating the statement of the rule. Basically, there are two possible situations. Either the sheets are connected together in the same fashion on either side of the link in question (the first three rows in Fig. 9), or they are connected together in a different fashion (the last three rows). By construction, the rule is invariant under permutation of the colours. So all you need to see is one of the first three rows and one of the last three, to reconstruct the whole table. There are, in each case, five possible choices for how to connect the sheets across the link in question. Of the five, the first three lead to the same value of $w_2(\Sigma)$, and the last two change $w_2(\Sigma)$ by $1^\star$. Thus, according to (3.5), they get counted with a minus sign relative to the first three. As in 2D, this leaves a *net* of one configuration which contributes.

This pretty much takes care of the proof that the sum (3.5) over unicursal surfaces constructed according to the rule (Fig. 9) reproduces exactly the Ising model partition function. The one point which should be clarified is the notion of nearest neighbour link. We have four sheets coming together along a curve. If the link under consideration is somewhere in the middle of that curve, then it is clear what nearest neighbour means: the next link along the curve. It may be, however, that the link in question is at the endpoint of the curve (as in Fig. 6). But then, it is also unambiguous which sheets are connected to which "to the left" of the link in question in Fig. 6. To illustrate this, Fig. 10

* If you have trouble seeing this, recall the definition of $\chi(\Sigma)$ as *vertices−edges+faces*, and count the *relative* contribution to $\chi(\Sigma)$ for each of the entries of a given row of Fig. 9. $\Delta\chi$ for the last two entries in a row relative to the first three is *odd*.



Fig. 9: Allowed configurations on one link



Fig. 10: Two configurations from the table above, and their interpretation

contains two examples of configurations from the second row of Fig. 9, where the "curve" in question is precisely one link in length. On the right, you can see the path of a "bug" on the corresponding surfaces.

There is one case where the meaning of nearest neighbour *is* still ambiguous, namely when three sheets intersect each other at a point. A link adjoining this vertex has five candidates for the role of "nearest neighbour". The simplest way to resolve this ambiguity is to define the nearest neighbour as the link straight ahead, since it is clearly singled out from the other four.

Having shown that the sum over unicursal surfaces reproduces the Ising partition function, one would like to take the next step in this program and relax the unicursal condition, by showing that the sum over admissible surfaces with weights given by (3.5) *equals* the sum over unicursal surfaces. I have looked at simple examples in which this seems to work, but I have not been able to construct a general proof along the lines of the one presented in the last section for the 2D case.

Assuming that the replacement of unicursal by admissible surfaces goes through, then the continuum limit of (3.5) is *obvious*. The sum over surfaces on the lattice becomes a functional integral over immersed surfaces, $A[\Sigma]$ simply becomes the Nambu-Goto action, and $w_2(\Sigma)$, being a topological-invariant, is perfectly well-defined, both in the continuum and on the lattice. Basically, we end up with the bosonic string in three dimensions – but with two crucial differences: 1) we sum over both oriented and non-oriented 2-surfaces, and 2) we weight surfaces with odd Euler characteristic with a *minus sign*.

4. Comments on quantization

When we recast the 2D Ising model as a sum over admissible curves, we were fortunate to find that the free energy $W[\beta]$ was given by a sum over *connected* curves *i.e.* over maps from the circle into the plane. Perhaps the most unattractive feature of recasting the 3D Ising model as a string theory is that even the sum over connected surfaces involves a sum over all genera of surface. This sum over genus is, in ordinary string perturbation theory, controlled by a (purportedly) small coupling constant g_{st}. In the Ising model, g_{st} has magnitude 1, since we want to count domain boundaries of all genus with equal weight. This is an unpleasant prospect to contemplate, since string perturbation theory is very likely a divergent series*.

We have somewhat improved this situation in the current formulation, since effectively we have taken $g_{st} = -1$. If the large-order behaviour of the perturbation series found in the matrix model [23] is any guide, this is enough to make the series Borel-summable$^\clubsuit$.

* Of course, in the low-temperature expansion, which has motivated our discussion, low-genus surfaces dominate because they tend to be the surfaces of minimal area. However, we are precisely interested in the theory near the critical point, where this area-suppression breaks down.

$^\clubsuit$ For oriented surfaces, the series goes like $(2g)!g_{st}^{(2g-2)}$. For non-oriented surfaces, one has also odd powers of g_{st}, and the series goes like $n!g_{st}^{(n-2)}$, which is Borel-summable for negative g_{st}.

But the summability of the perturbation series is a rather distant concern compared with the more immediate problem of the tachyon divergence of *each term* in the series. Here, too, introducing a negative string coupling introduces a measure of hope. Imagine introducing, in some uniform way, a cutoff on the modular integration for each genus. This renders each term in the sum finite. The contribution from each genus diverges as we send the cutoff to zero, but now, because there are explicit minus signs in the alternating sum, these divergences have the possibility of cancelling *between* different genera. Effectively, when we pinch a cycle as in Fig. 11, we could be pinching an orientable or a non-orientable surface. These contribute with the same weight, but opposite sign, to the sum over surfaces. If our cutoff is sufficiently clever, *all* of the divergent pieces may cancel, leaving a result which is *finite* as we take the cutoff to zero. This may seem like outrageous handwaving, but the lattice theory is an explicit realization of just such a cutoff. Hopefully, one can find a more convenient one for the purposes of continuum calculations.



Fig. 11: A pinched surface could be the limit of an orientable or a non-orientable surface, depending on the nature of the cycle(s) pinched.

Not only do the divergent terms cancel, which is to say that the contribution of the tachyon in the pinched channel cancels out, but it is reasonable that the contribution of many other states of the string cancel as well. In this way, the strongly-coupled string theory may have many fewer propagating states than we expected. Indeed, it is only through such a drastic set of cancellation that a string theory could ever reproduce the critical behaviour of the Ising model, which is, after all, that of a field theory.

This mechanism, for cancelling divergences and reducing the number of propagating states of a string theory is certainly novel. If, indeed, it works (as it must, if the Ising model is to be described as a string theory), then it is a telling example of just how different strongly-coupled string dynamics can be from that seen in perturbation theory. The details, of course, are peculiar to D=3, and $|g_{st}| = 1$. After all, the form of the string integrand is D-dependent, and if this cancellation is to be realized for D=3, it will, in general, not take place for other values of D. Similarly, the string coupling constant is

necessarily fixed by demanding that this cancellation take place. Hence we expect that the dilaton, too, is projected out of the spectrum[†].

With this vast truncation of the spectrum, string theory is clearly not a very *economical* description of the long-distance physics of the Ising model. But then again, it is not a very economical description of elementary particle physics either (which is also well-described as a field theory). The key question we as string theorists are grappling with is how string theory with its good short-distance behaviour can describe the long-distance physics that we see. The example of the Ising model should caution us that the answer may be stranger than we think.

Appendix A. Undoubled lattice fermions, and the 2D Ising model

As mentioned in the main text, the continuum limit of the 2D Ising model is a free majorana fermion. This is most easily seen by formulating it as a theory of lattice fermions, which is the subject of this appendix. When one does so, one immediately is confronted with the problem of fermion doubling[⋆]. Fermion doubling is a very generic feature of lattice fermion theories. Should we be surprised that the theory of lattice fermions had as its continuum limit a *single* majorana fermion which is massless at the critical point? How does this theory manage to avoid a doubling of the spectrum, i.e. why does the continuum limit not produce two (or four) species of free majorana fermions?

If it did produce four majorana fermions, then the central charge, as measured by the finite size scaling [24] would be $c = 2$, rather than $c = 1/2$. Other critical exponents (such as the divergence of the correlation length as we approach the critical point) would be the same for four decoupled Ising models as for one, but the discrepancy in c rules out this possibility.

We will examine this question both for the Ising model and for a closely related model, the Ashkin-Teller model.

Let us begin with the Ising model. Following [6], one can introduce a theory of lattice fermions whose partition sum reproduces exactly the diagrammatic expansion (2.3) of the

[†] The dilaton had better be projected out for another reason: were it not projected out, its presence would still lead to a divergence in Fig. 11.

[⋆] I'd like to thank P. Ginsparg for sharing his insights into this subject

2D Ising model. Introduce fermionic variable $\psi_1, \psi_2, \psi_3, \psi_4$ living on the sites of the dual lattice with the action

$$\begin{aligned} S = \sum_{\mathbf{x}} \ & \mathrm{e}^{-2\beta}(\psi_1 \hat{\partial}_x \psi_3 + \psi_2 \hat{\partial}_y \psi_4) + \psi_1\psi_2 + \psi_2\psi_3 + \psi_3\psi_4 + \psi_1\psi_4 \\ & + (1 + \mathrm{e}^{-2\beta})(\psi_1\psi_3 + \psi_2\psi_4) \end{aligned} \tag{A.1}$$

($\hat{\partial}_x \psi = \psi(\mathbf{x} + \hat{\imath}) - \psi(\mathbf{x})$ is the lattice derivative.) The expansion of $\int [D\psi] \mathrm{e}^S$ reproduces the low-temperature expansion (2.3). To see this, note that we need to bring down four fermions per site in order to get a nonzero grassmann integral. We can do this in one of two ways: either by bringing down "propagator terms", $\psi_1(\mathbf{x})\psi_3(\mathbf{x}+\hat{\imath})$ or $\psi_2(\mathbf{x})\psi_4(\mathbf{x}+\hat{\jmath})$, each of which comes with a factor of $\mathrm{e}^{-2\beta}$, or by bringing down "mass terms" from (A.1). The former must form closed curves, which get weighted by $\mathrm{e}^{-2\beta L}$. Sites which are not visited by these curves must have two mass terms brought down from the action. There are three different ways of doing this, but by Fermi statistics, one gets counted with a minus sign. When the dust settles, we precisely reproduce the low-temperature expansion (2.3).

In the continuum limit (A.1) reduces to the action for a free majorana fermion of mass $m = (\mathrm{e}^{2(\beta-\beta_c)} - 1)/a$ (a is lattice spacing). At the critical point, the fermion is massless. To study the continuum limit, it is helpful to form new linear combination of the fermions which diagonalize the mass term in (A.1) [25]. Let $\alpha = \mathrm{e}^{i\pi/4}$ and form the linear combinations

$$\begin{aligned} \psi &= \tfrac{1}{2}[\alpha^7\psi_1 + \alpha^2\psi_2 + \alpha^5\psi_3 + \psi_4] \\ \tilde{\psi} &= \tfrac{1}{2}[\alpha\psi_1 + \alpha^6\psi_2 + \alpha^3\psi_3 + \psi_4] \\ \chi &= \tfrac{1}{2}[\alpha^3\psi_1 + \alpha^2\psi_2 + \alpha\psi_3 + \psi_4] \\ \tilde{\chi} &= \tfrac{1}{2}[\alpha^5\psi_1 + \alpha^6\psi_2 + \alpha^7\psi_3 + \psi_4] \quad . \end{aligned}$$

The action (A.1) becomes

$$\begin{aligned} S = \sum_{\mathbf{x}} \ \mathrm{e}^{-2\beta} \Big[& -\frac{1}{4}(\psi-\chi)\hat{\partial}_x(\psi-\chi) - \frac{1}{4}(\tilde{\psi}-\tilde{\chi})\hat{\partial}_x(\tilde{\psi}-\tilde{\chi}) \\ & - \frac{i}{4}(\psi+\chi)\hat{\partial}_y(\psi+\chi) + \frac{i}{4}(\tilde{\psi}+\tilde{\chi})\hat{\partial}_y(\tilde{\psi}+\tilde{\chi}) \\ & + \frac{i}{4}\big(\hat{\partial}_x(\psi-\chi)\hat{\partial}_x(\tilde{\psi}-\tilde{\chi}) + \hat{\partial}_y(\psi+\chi)\hat{\partial}_y(\tilde{\psi}+\tilde{\chi})\big) \\ & + i\hat{m}_1\psi\tilde{\psi} - i\hat{m}_2\chi\tilde{\chi}\Big] \end{aligned} \tag{A.2}$$

where

$$\hat{m}_1 = (\mathrm{e}^{2(\beta-\beta_c)} - 1), \qquad \hat{m}_2 = 2\sqrt{2}\mathrm{e}^{2\beta} + \hat{m}_1, \qquad \mathrm{e}^{2\beta_c} = 1 + \sqrt{2} \quad . \tag{A.3}$$

The continuum limit is obtained by letting $\beta \to \beta_c$, while the lattice spacing $a \to 0$, holding $m_1 = \hat{m}_1/a$ fixed. In this limit, $m_2 = \hat{m}_2/a \sim 2\sqrt{2}/a \to \infty$, so we can simply drop all dependence on $\chi, \tilde{\chi}$. Looking at the kinetic energy operator that remains for $\psi, \tilde{\psi}$, one notices two important features. First, because it is defined using *asymmetric* lattice derivatives, it is non-Hermitian. In momentum space $\hat{\partial}_x \psi = \psi(\mathbf{x} + \hat{\imath}) - \psi(\mathbf{x}) \to 2ie^{ip_x/2} \sin(p_x/2)\psi(\mathbf{p})$, unlike the symmetric lattice derivative $\frac{1}{2}(\psi(\mathbf{x} + \hat{\imath}) - \psi(\mathbf{x} - \hat{\imath})) \to i \sin(p_x)\psi(\mathbf{p})$. Second, the kinetic energy operator has only one zero in the Brillouin zone, at $\mathbf{p} = 0$. The latter feature is, of course, what we want. The former is a little troubling. Generally, it is believed that a non-Hermitian kinetic energy operator would lead to non-unitarity of the continuum theory. Here it is clearly harmless. Indeed, expanding about $\mathbf{p} = 0$, one simply replaces lattice derivatives by continuum derivatives $\hat{\partial} \to a\partial$ and, after rescaling the fields, one obtains

$$S_{cont} = \int \mathrm{d}^2 z \left(-\tfrac{1}{2}\psi\partial_{\bar{z}}\psi - \tfrac{1}{2}\tilde{\psi}\partial_z\tilde{\psi} + im_1\psi\tilde{\psi}\right) \tag{A.4}$$

which is, as promised, the action for a free majorana fermion of mass m_1 ($\partial_{z,\bar{z}} = \frac{1}{2}(\partial_x \mp i\partial_y)$). All of the non-Hermiticity of the lattice action went into *irrelevant* operators[†] which can be neglected in the continuum limit.

Even global effects are captured correctly. Recall from our discussion of [5], that for the fermions to reproduce the Ising partition function on a lattice with periodic boundary conditions, we need to introduce independent horizontal and vertical couplings and sum over the partition functions where we shift the horizontal and vertical couplings by $i\pi/2$

$$\mathcal{Z}[\beta] = \tfrac{1}{2}[Z(\beta,\beta) + Z(\beta + i\pi/2, \beta) + Z(\beta, \beta + i\pi/2) + Z(\beta + i\pi/2, \beta + i\pi/2)]$$

These shifted partition functions are generated by the modified fermionic actions

$$\begin{aligned} S = \sum_{\mathbf{x}} \; & \mathrm{e}^{-2\beta}(\pm\psi_1\hat{\partial}_x\psi_3 \pm \psi_2\hat{\partial}_y\psi_4) + \psi_1\psi_2 + \psi_2\psi_3 + \psi_3\psi_4 + \psi_1\psi_4 \\ & + (1 \pm \mathrm{e}^{-2\beta})\psi_1\psi_3 + (1 \pm \mathrm{e}^{-2\beta})\psi_2\psi_4 \end{aligned} \tag{A.5}$$

[†] The fact that Wilson-like terms remove the doubling in the Ising model was noted previously by Hassenfratz and Maggiore [26].

Going through the same analysis as before, one sees that modifying the action in this way changes the boundary conditions for the continuum fermion from periodic to antiperiodic, so that (2.7)indeed reproduces the sum over spin structures for the fermion. All of these somewhat subtle features of the continuum partition function are reproduced by the lattice theory without doubling or other inconsistencies.

One might think that everything worked out so nicely here because the continuum limit is a *free* theory. This is not so. Consider instead the Ashkin-Teller model. This is a theory of two Ising models coupled by a four-spin interaction. In the fermionic formulation, it consists of two copies of (A.1) (for $\psi_{1,\ldots,4}$ and $\psi_{5,\ldots,8}$), coupled by a four-fermi interaction

$$\Delta S = \kappa\big((\psi_1\hat{\partial}_x\psi_3 + \psi_1\psi_3)(\psi_5\hat{\partial}_x\psi_7 + \psi_5\psi_7) + (\psi_2\hat{\partial}_y\psi_4 + \psi_2\psi_4)(\psi_6\hat{\partial}_y\psi_8 + \psi_6\psi_8)\big)$$

In the continuum limit, this reduces to a massive Thirring model♣ (massless along the critical line). This is a genuine interacting field theory. Interestingly, the *location* of the critical point (line) is not shifted by the interactions or, to say it differently, the ultra-massive doublers (and the χ's) do not feed down their masses to the "light" fermions through the interactions. The light fermion masses are protected, not through a continuous chiral symmetry, but through a discrete symmetry – the duality symmetry of the Ashkin-Teller model. This use of a duality symmetry to keep light fermions light is a somewhat novel mechanism from the point of view of lattice field theory. Perhaps it has some wider applicability.

Appendix B. Is Nambu-Goto = Polyakov ?

Most of the comments made in Section 4, tacitly assumed that one knew, at least in principle, how to quantize the string theory. For the Nambu-Goto string, we really know very little, but it is usually assumed that Nambu-Goto is equivalent to the Polyakov string.

The usual argument [10] proceeds as follows. Introduce an intrinsic metric $g = f^*(\mathrm{e}^{\phi}\hat{g})$ on the surface Σ, where $\hat{g}$ is a fiducial metric and f a diffeomorphism. Rewrite the Nambu-Goto action as

$$\int \sqrt{g} + i\lambda^{ab}(\partial_a\mathbf{X}\cdot\partial_b\mathbf{X} - g_{ab}) \tag{B.1}$$

♣ Perhaps, I should say "Thirring-like" model. It consists of two massive majorana fermions coupled by a four-fermion interaction $\lambda\int \mathrm{d}^2z\ \psi\tilde{\psi}\psi'\tilde{\psi}'$.

where λ^{ab} is a lagrange-multiplier field which sets the intrinsic metric g_{ab} equal to the induced metric $h_{ab} = \partial_a \mathbf{X} \cdot \partial_b \mathbf{X}$. (I have written the intrinsic metric in the conformal gauge because that's where I understand best how to quantize the theory, but the following remarks hold true in the light-cone gauge as well.)

If we assume that the lagrange-multiplier field λ has short-range correlations, then one can solve for it using the equations of motion, in which case, (B.1) becomes equal to the Polyakov action.

Unfortunately, for a 3-dimensional target space, the generic singularity of the induced metric h has *one* of its eigenvalues going to zero, while the other remains finite. We would not be too upset if both of the eigenvalues went to zero (as happens for a 1-dimensional target space); that would simply correspond to $\phi \to -\infty$, which is simply the boundary of field-space and is "included" in the ϕ functional integral. But this Whitney-type singularity [27], where one eigenvalue goes to zero, does not correspond to *any* configuration of the field ϕ. It is simply inaccessible in conformal gauge. (More properly, it is related by a *singular* "diffeomorphism" to a metric in the conformal gauge slice.)*

This should make one a little worried (but perhaps only a little) about the equivalence of between the Nambu and Polyakov actions for D=2,3, where the generic singularity is of the Whitney type.

Appendix C. Fermions on nonorientable manifolds

Because of the difficulties alluded to in the text, it must be the case that the definition of fermions used in open string theory *must* be different from that studied by Grinstein and Rohm [19]. In particular, the mœbius band, viewed in the closed string channel, has both a NS-NS sector and an R-R sector, whereas the fermions discussed in [19] have only an R-R sector. Here I will explain the discrepancy: there are two *inequivalent* definitions of fermions on a nonorientable manifold.

The pinor bundle has a structure group which is a $\mathbb{Z}_2$ extension of the structure group of the manifold

$$0 \to \mathbb{Z}_2 \to Pin(n) \to O(n) \to 0 \tag{C.1}$$

* Sedrakyan[28] has argued that one might try to mimic the effect of these Whitney singularities by a gas of operator insertions on an otherwise smooth surface. In the most naïve approach, operator insertions simply produce ordinary conical singularities, which is not what we want.

The mœbius band and the Klein bottle (the only nonorientable manifolds for which explicit calculations have been done by open string theorists) are both flat 2-manifolds. Thus the structure group can be reduced from $O(2)$ to $\mathbb{Z}_2$. There are clearly two possible $\mathbb{Z}_2$ extensions of the reduced structure group: $\mathbb{Z}_4$ or $\mathbb{Z}_2 \times \mathbb{Z}_2$. The definition of fermions used by open string theorists corresponds to the former (the transition functions of the fermions are in $\mathbb{Z}_4$), whereas the fermions in [19] correspond to the latter extension.

It is not immediately evident whether the former definition can be extended to curved 2-manifolds with the full $O(2)$ structure group. As it turns out, it can [22]*. Here I will just summarize the results, refering to [22] for details.

One starts with the Clifford algebra $\mathcal{C}_n^\pm$ (over $\mathbb{R}$), generated by the Dirac matrices γ_a with the relation

$$\gamma_a\gamma_b + \gamma_b\gamma_a = \pm 2\delta_{ab}\mathbb{I} \tag{C.2}$$

On $\mathcal{C}_n^\pm$, one defined a "norm" by

$$\|\gamma_{v_1} \dots \gamma_{v_N}\|^2 = |\mathrm{Tr} u^2|$$

(where Tr is the usual quadratic form on the Clifford algebra, normalized so that $\mathrm{Tr}\mathbb{I} = 1$) and the group $Pin^\pm(n)$ is the multiplicative group of invertible elements of $\mathcal{C}_n^\pm$ of unit norm. In each case, $Pin(n)$ is the extension by $\mathbb{Z}_2$ of $O(n)$, but in general $Pin^+(n) \not\simeq Pin^-(n)$. In particular, consider $Pin(1)$, which consists of 4 elements, $\{\pm\mathbb{I}, \pm\gamma\}$. But depending on whether $\gamma^2 = \pm\mathbb{I}$, we obtain different groups: $Pin^+(1) = \mathbb{Z}_2 \times \mathbb{Z}_2$, while $Pin^-(1) = \mathbb{Z}_4$.

$Spin^\pm(n)$ is the subgroup of $Pin^\pm(n)$ which extends the identity component of $O(n)$

$$0 \to \mathbb{Z}_2 \to Spin^\pm(n) \to SO(n) \to 0$$

However, one always finds that $Spin^+(n) \simeq Spin^-(n)$. Thus *for orientable manifolds* the choice of sign in (C.2) turns out to be irrelevant to defining spinors. However, for nonorientable manifolds, *it matters*!

We then go on to try to use the group extension (C.1) to lift the transition functions of the $O(n)$ frame bundle on M to those of a principal $Pin^\pm(n)$ bundle. In so doing, we encounter an obstruction. Even though the $O(n)$ transition functions obey the cocycle condition

$$g_{ij} \cdot g_{jk} \cdot g_{ki} = \mathbb{I} \quad ,$$

* I'd like to than R. Rohm for bringing this reference to my attention.

when lifted to $Pin(n)$ transition functions, they may not:

$$\tilde{g}_{ij} \cdot \tilde{g}_{jk} \cdot \tilde{g}_{ki} \in \mathbb{Z}_2 \quad .$$

This defines an element of $\mathrm{H}^2(M, \mathbb{Z}_2)$. For $Pin^+(n)$, the obstruction is the second Stiefel-Whitney class $w_2(M)$. For $Pin^-(n)$, the obstruction is $w_2 + w_1 \cup w_1$. The former, as noted in [19], is nonvanishing on certain 2-manifolds. Hence one cannot define a Pin^+ bundle on those manifolds. The latter (as one sees from equations (3.3),(3.4)) *vanishes identically* on any 2-manifold. Thus, we can always define a Pin^- structure on any 2-manifold and this (not the more intuitive Pin^+) is the definition of fermions used by open string theorists[†].

Acknowledgments

I would like to thank M. Bershadsky, P. Ginsparg, D. Kutasov, A. Polyakov, and H. Verlinde for useful conversations. I would especially like to thank A. Sedrakyan, conversations with whom kindled my interest in this subject. This work was supported by NSF grant PHY90-21984.

[†] The $SO(32)$ open string defined using Pin^+ is a rather amusing variant. It is an anomaly-free string theory with a nonvanishing cosmological constant (and hence a dilaton tadpole). In that respect, it resembles the $SO(16) \times SO(16)$ heterotic string. The theory is anomaly free because the mœbius band and cylinder amplitudes in the R-R sector are exactly the same as in the usual $SO(32)$ open string. However the dilaton tadpole does not cancel between the disk and $\mathbb{R}P^2$, as it does in the usual $SO(32)$ string. Rather, the dilaton 1-point function on $\mathbb{R}P^2$ – defined physically by sewing – vanishes identically.

References

[1] L. Onsager, Phys. Rev. **65** (1944) 177.

[2] M. Kac and J. C. Ward, Phys. Rev. 88 (1952) 1332.

[3] R. Feynman, *Statistical Mechanics, A Set of Lectures*, W. A. Benjamin (1972).

[4] S. Sherman, J. Math. Phys. **1** (1963) 202, *ibid* **4** (1963) 1213.

[5] R. Potts and J. C. Ward, Prog. Theor. Phys. **13** (1955) 38.

[6] C. Hurst and H. Green, J. Chem. Phys. **33** (1960) 1059.

[7] T. Schultz, D. Mattis and E. Lieb, Rev. Mod. Phys. **36** (1964) 856.

[8] E. Fradkin, M. Srednicki and L. Susskind, Phys. Rev. **D21** (1980) 2885.

[9] A. Polyakov, Phys. Lett. **103B** (1981) 211.

[10] A. Polyakov, *Gauge Fields and Strings*, Harwood Academic Publishers (1987).

[11] Vl. Dotsenko and A. Polyakov, *in* Advanced Studies in Pure Math 15;
Vl. Dotsenko, Nucl. Phys. **B285** (1987) 45.

[12] A. Casher, D. Fœrster and P. Windey, Nucl. Phys. **B251** (1985) 29;
C. Itzykson, Nucl. Phys. **B210** (1982) 477.

[13] A. Sedrakyan, Phys. Lett.. **137B** (1984) 397;
A. Kavalov and A. Sedrakyan, Phys. Lett. **173B** (1986) 449;
A. Kavalov and A. Sedrakyan, Nucl. Phys. **B285** (1987) 264.

[14] D. Kutasov and N. Seiberg, Nucl. Phys. **B358** (1991) 600.

[15] L. Alvarez-Gaumé, G. Moore and C. Vafa, Comm. Math Phys. **106** (1986) 40.

[16] For a discussion, see T. Jarosewicz and P. Kurzepa, Ann. Phys. **210** (1991) 255.

[17] E. Moise, *Geometric Topology in Dimensions 2 and 3*, Springer-Verlag (1977)

[18] J. Milnor and J. Stasheff, *Characteristic Classes*, Princeton University Press (1974)

[19] B. Grinstein and R. Rohm, Comm. Math. Phys. 106 (1986) 40.

[20] M. Atiyah, R. Bott and A. Shapiro, Topology **3** (1964) 3.

[21] M. Green and J. Schwarz, Phys. Lett. **151B** (1985) 21.

[22] Y. Choquet-Bruhat and C. DeWitt-Morette, "*Analysis, Manifolds and Physics*, part II: Applications", North Holland (1989).

[23] P. Ginsparg and J. Zinn-Justin, Phys. Lett.. **B255** (1991) 189;
S. Shenker, *in* 1990 Cargèse Workshop on Random Surfaces, Quantum Gravity and Strings, RU90-47.

[24] H. Blöte, J. Cardy , and M. Nightingale, Phys. Rev. Lett.. **56** (1986) 742;
I. Affleck, Phys. Rev. Lett.. **56** (1986) 746.

[25] V. Dotsenko and Vl. Dotsenko, Adv. Phys. **32** (1983) 129.

[26] P. Hassenfratz and M. Maggiore, Phys. Lett. **B220** (1989) 215.

[27] M. Golubitsky and V. Guillemin, *Stable Mappings and Their Singularities*, Springer-Verlag (1973).

[28] A. Sedrakyan, talk at Princeton, Fall, 1991.